

Nonlinear Inversion from Nonlinear Filters for Ocean Acoustics

Robert I. Odom
Applied Physics Laboratory
College of Ocean and Fisheries Sciences
University of Washington
1013 N.E. 40th Street
phone: (206)685-3788 fax: (206)543-6785 email: odom@apl.washington.edu

Award # N00014-05-1-0087

LONG TERM GOALS

The long term goals of this research are to develop practical and efficient algorithms for application the nonlinear inversion problems encountered in ocean acoustics. Such algorithms would be used for estimating or accounting for the effects of the environment on acoustic propagation, detection and tracking in shallow water.

OBJECTIVES

The specific objectives of this research are adapt a specific nonlinear filter, known as a Daum filter, for acoustic inversion of shallow water environmental properties, and to assess the performance of this nonlinear filter relative to local linear inversion on the one hand and global methods, e.g. Monte Carl methods on the other hand.

APPROACH

Many inverse problems of interest in ocean acoustics are intrinsically nonlinear, e.g. inverting measured pressure data for bottom and scattering properties. The solution to the nonlinear inversion problem is usually approached in one of two ways. The first way is to assume a starting model, which one hopes is near to the true model, then recursively solve a linearized version of the inverse problem for corrections to the starting model and model covariance. The advantage of this approach is that the numerical implementation of the solution algorithm is relatively straightforward and in a linear problem the statistical properties are well defined and will remain gaussian if they start out gaussian. However linearization of a nonlinear system can produce biased estimates for two reasons: 1. Linearization of the system and/or measurement equations may not be a good approximation, and 2. Nonlinear systems do not maintain gaussian statistics as they evolve even if they are initially gaussian. Another problem with linearizing a nonlinear system is that with a poor starting guess the solution algorithm may never converge to the true answer. If the starting model represents a point near a local minimum of the solution space, the final solution will be trapped in that local minimum, and never converge to the true answer. This can be circumvented by using Monte Carlo techniques to randomly sample the solution space for starting models.

The other class of solution methods attack the nonlinear problem directly by using simulated annealing or genetic algorithms. The disadvantage of these directly nonlinear methods, is that there is

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2005		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE Nonlinear Inversion from Nonlinear Filters for Ocean Acoustics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, College of Ocean and Fisheries Sciences, 1013 N.E. 40th Street, Seattle, WA, 98195				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT The long term goals of this research are to develop practical and efficient algorithms for application the nonlinear inversion problems encountered in ocean acoustics. Such algorithms would be used for estimating or accounting for the effects of the environment on acoustic propagation, detection and tracking in shallow water.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

1. The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that proper record-keeping is essential for the integrity of the financial system and for the ability to detect and prevent fraud.

2. The second part of the document outlines the specific procedures for recording transactions. It details the steps involved in the accounting cycle, from identifying the transaction to posting it to the appropriate ledger account.

3. The third part of the document discusses the role of the auditor in verifying the accuracy of the records. It describes the various techniques used by auditors to test the reliability of the data and to ensure that the financial statements are presented fairly.

4. The fourth part of the document addresses the issue of internal controls. It explains how a well-designed system of internal controls can help to minimize the risk of error and to ensure that the organization's assets are protected.

5. The fifth part of the document discusses the importance of transparency and accountability in financial reporting. It argues that organizations should be open and honest about their financial performance and should provide clear and concise information to their stakeholders.

6. The sixth part of the document discusses the role of the government in regulating the financial system. It describes the various laws and regulations that govern the behavior of financial institutions and the consequences of non-compliance.

7. The seventh part of the document discusses the importance of ethical behavior in the financial industry. It argues that financial professionals should always act in the best interests of their clients and should avoid any conflicts of interest.

8. The eighth part of the document discusses the importance of ongoing education and training for financial professionals. It argues that the financial industry is constantly evolving and that professionals must stay up-to-date on the latest developments.

9. The ninth part of the document discusses the importance of collaboration and communication between financial institutions and regulators. It argues that a strong working relationship is essential for the effective supervision of the financial system.

10. The tenth part of the document discusses the importance of public confidence in the financial system. It argues that the financial system can only function properly if the public has faith in its integrity and in the actions of its regulators.

no way to conveniently propagate the statistical properties of the solution through to the final result. One solution to this problem is to find the global minimum in the solution space, if one exists, then linearize about the solution representing the global minimum and do a statistical analysis about that solution. This was done by Potty et al.(2000), who employed a genetic algorithm followed by linear analysis about the solution determined by the genetic algorithm.

The recursive algorithms commonly employed for the estimation of the model and covariance relative to some initial starting values bear a strong resemblance to Kalman filters, which are commonly employed in target tracking algorithms. The original Kalman filter was derived for strictly linear systems. However, the Extended Kalman Filter can be applied to systems which are weakly nonlinear. In the late 1980s Frederick Daum, a mathematician working at Raytheon Corporation, developed a fully nonlinear formulation to the filtering problem for target tracking (Daum, 1985, 1986, 1987). His theory is elegant, but impractical from an implementation point of view. Sometime later Schmidt (Schmidt, 1993) succeeded in deriving an approximate algorithm based on Daum's original theory, and developed a successful numerical implementation of a nonlinear filter that was a significant improvement to the Kalman and Extended Kalman filters for the type of tracking problem Schmidt was interested in.

This research aims to develop an ocean acoustic inversion algorithm based on Schmidt's (1993) implementation of Daum's nonlinear filtering theory. The purpose is to be able to carry along the statistics of the geoacoustic model parameters through the inversion process. The work is much more than a straightforward "rename the variables and code it up". However, the tracking algorithms do bear resemblance to the iterative inversion algorithms for updating model parameter means and covariances of an iterated inverse problem as in, for example, Menke (1983). Most estimation problems can be cast into an iterative form, whereby the state vector, which in our case is the ocean bottom model vector, is updated sequentially as data is added. Estimation filter formulations are also natural for range dependent or time dependent environments. Daum's original theory and Schmidt's practical implementation assumes nonlinear dynamics and a linear relationship between the measurement and state vectors. In our case the measurement vector, complex pressure say, and the state vector, the bottom model, are not linearly related. The filter needs to be re-derived from scratch with the measurement to state vector relationship appropriate for our ocean acoustic application. Once re-derived, it will need to be coded, and checked against results for linear inverse problems. Dosso (e.g. Dosso and Wilmut, 2002) at the University of Victoria has developed a Monte Carlo inversion method for the ocean acoustics problem. This is computationally very intensive, but he does get the full probability density function (pdf) for the model parameters. Because the Schmidt implementation of the Daum theory propagates the additional terms in the mean and covariance of the state vector pdf, it falls in between the standard linear inversion methods and Dosso's Bayesian Monte Carlo methods.

Currently employed algorithms for nonlinear problems such as simulated annealing and genetic algorithms have no mechanism for propagating the statistics. What the nonlinear filter algorithm will do is provide a natural mechanism for updating the statistics as a solution is determined. A comparison of the filter with an algorithm such as simulated annealing would be illuminating, and a valuable check on the filter algorithm itself.

Filter type algorithms are ideally suited to inverse problems with time dependent oceanography or range dependence. We do not anticipate attempting to include time dependent oceanography at this time, but we would like to look at the issue of range dependent inversion. The idea would be to sequentially update parameter estimates as a function of range. Also note that any inversion algorithm

can be cast into a filter like algorithm by supplying the data sequentially and updating the model parameter estimates sequentially as data is added to the problem, or a smoother by considering the complete data set, and working both forwards and backwards through the data set. In the end, probably the best formulation to use for a given inverse problem depends on the noise statistics. This is also something we propose to investigate.

Linear inverse problems admit the construction of both data and model resolution matrices. These resolution matrices can be used as metrics with which to estimate model uniqueness and data predictability. We will be able to construct resolution matrices for the nonlinear problem and compare them with their fully linear equivalents.

WORK COMPLETED

Previously we have taken a deterministic view of the resolution matrix. When noise or incomplete measurements are included, a statistical view must be adopted. The well known Fisher information matrix is directly related to the model resolution matrix. The Fisher information matrix is $\mathbf{F} = \mathbf{G}^g \mathbf{E}^{-1} \mathbf{G}$, and the resolution matrix is $\mathbf{R} = \mathbf{A} \mathbf{F}$, where \mathbf{E} is the covariance of the measurements plus the error in the forward model, \mathbf{A} is the covariance of the model estimate after incorporation of the measurement – the *a posteriori* covariance, \mathbf{G} is the matrix of model sensitivities, the partial derivatives w.r.t. the model parameters, and \mathbf{G}^g denotes the generalized inverse of \mathbf{G} . The trace of the resolution matrix is simply related to the total number of model parameters of the *a priori* and *a posteriori* covariances. In the next work period will be exploring these relationships numerically and developing relations for a nonlinear version of the resolution matrix.

As stated above an inverse problem can be recast as a filtering problem. A strictly linear problem becomes a Kalman Filter (KF). A problem that has been locally linearized becomes an Extended Kalman Filter (EKF), and the fully nonlinear problem can be represented as a Daum – Schmidt Filter. We have completed working computer code for a simple EKF with which to compare more complicated inverse models.

RESULTS

Figure 1 shows numerical results for the comparison of an EKF (white) and gradient search (black) applied to a weakly nonlinear test problem. The location of a stationary source is solved for using 10 receivers each with 10 noisy data points. Each method starts from some initial guess and has a 95% confidence ellipse about its final point. The colorful surface is the objective surface. The red square, which is the solution point, is not exactly at the minimum of the surface. The data have additive Gaussian noise, which slightly biases the surface. Regularization of the gradient descent was done by truncating the small eigenvalues. We developed the EKF code as this will be the algorithm against which we will compare the Daum Filter.

Figure 2 shows the convergence of the location as more receivers, each with 10 data points are incorporated into the source location estimate.




Figure 1. *Comparison of gradient descent (black) with an EKF (white) for a slightly nonlinear source localization problem. The yellow triangles are the receiver locations, each with 10 data points. Noise in the measurements causes the bias in the final location.*




Figure 2. *Model error for the example in Figure1. As more receivers, each with 10 data points with noise, are included, the final source location is estimated.*

IMPACT/APPLICATIONS

A nonlinear, well characterized filter-based inversion method and algorithm will have application to environmental estimation and target tracking. A practical method way to compute the resolution for a nonlinear inversion will have an impact on the characterization of uncertainty and uniqueness of environmental estimates required for acoustic propagation.

RELATED PROJECTS

Our research is directly related to other programs studying effects of uncertainty in the environment, measurements, and models on acoustic propagation, and target detection and characterization.

REFERENCES

Daum, F.E., "A New Nonlinear Filtering Formula for Discrete Time Measurements," *Proc. 24th IEEE Conf. Decision and Control*, Vol. **3**, pp1957-1958, Dec. 11-13, Fort Lauderdale, Florida, 1985.

Daum, F.E., "Exact Finite-Dimensional Nonlinear Filters," *IEEE Trans. Automatic Control*, **AC-31**, pp616-622, 1986.

Daum, F.E., "Application of New Nonlinear Filtering Theory," *Proc. 1987 Am. Control Conf.*, Vol. **3**, paper TA-1, Jun. 10-12, Minneapolis, Minnesota, 1987.

Dosso, S.E. and M.J. Wilmut, "Quantifying data information content in geoacoustic inversion", *IEEE J. Ocean. Eng.*, **27**, pp296-304, 2002.

Menke, W., *Geophysical data analysis: discrete inverse theory*, Rev. ed., 285 p., Academic Press, San Diego, California, 1989.

Potty, G., J.H. Miller, J.F. Lynch, K.B. Smith, "Tomographic inversion for sediment parameters in shallow water," *J. Acoust. Soc. Am.*, **108**, pp973-986, 2000.

Schmidt, G.C., "Designing Nonlinear Filters Based on Daum's Theory," *J. Guid. Control, and Dynamics*, **16**, pp371-376, 1993.

PUBLICATIONS

Ganse, A.A.. and R.I. Odom, "Adapting a truly nonlinear filter to the ocean acoustic inverse problem," (abstract), *J. Acoust. Soc. Am.*, **117**, p2463, 2005.

Odom, R.I., and A.A. Ganse, "Assessing uncertainty in ocean acoustic inverse problems," (Invited talk and abstract), *J. Acoust. Soc. Am.*, **118**, p1903, 2005.